



King's Research Portal

Document Version
Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Palmer, K. L., Law, J., Carter, B., Hewitt, J., Boyle, J., Maitra, C., Farrell, I., & Moug, S. (2019). Frailty in Older Patients undergoing Emergency Laparotomy: results from the observational ELF Study (Emergency Laparotomy and Frailty). *Annals of Surgery*, 48(Supplement_2), ii28–ii29. [98].

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**Frailty in Older Patients undergoing Emergency Laparotomy: Results from the U.K. observational
Emergency Laparotomy and Frailty (ELF) Study.**

Kat L Parmar, Jennifer Law, Ben Carter, Jonathan Hewitt, Jemma M Boyle, Patrick Casey,
Ishaan Maitra, Ian S Farrell, Lyndsay Pearce and Susan J Moug* on behalf of the ELF Study
Group.

Kat L Parmar: BSc, MBChB, MSc, PGCE, FRCSEng. Higher Surgical Trainee and Clinical Research Fellow, Manchester Cancer Research Centre, Wilmslow Road, Manchester M20 4GJ. klparmar@doctors.org.uk

Jennifer Law: BMBS, BMedSci, MSc. Higher Surgical Trainee and Clinical Research Fellow, University of Liverpool, Sherrington Building, Ashton Street, L69 3BX. jlaw1@nhs.net

Ben Carter: PhD, MSc. Senior Lecturer in Biostatistics Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology & Neuroscience, King's College London, De Crespigny Park, London, UK, SE5 8AF. ben.carter@kcl.ac.uk

Jonathan Hewitt: MB BS, MSc, FRCPGlas PhD. Clinical Senior Lecturer, Department of Population Medicine, 5th floor, NeuaddMeirionnydd, Heath Park Campus, Cardiff University. CF64 2XX. Hewittj2@cardiff.ac.uk

Jemma M Boyle: MB ChB, PGCE, MRCS. Higher Surgical Trainee and Clinical Research Fellow, Royal College of Surgeons of England, 35-43 Lincoln's Inn Fields, London, WC2A 3PE. jboyle@rcseng.ac.uk

Patrick Casey: MB ChB, MRCS. Higher Surgical Trainee, Health Education North West, Three Piccadilly Place, Manchester, M1 3BN. Patrick.casey1@nhs.net

Ishaan Maitra: BSc (Hons), MB ChB, MRCS. Higher Surgical Trainee and Research Registrar, NorthWest Deanery, Lancashire Teaching Hospitals NHS Foundation Trust, Preston, PR2 9HT. ishaan.maitra@googlemail.com

Ian S Farrell: MB ChB, MPharm, MRCS. Higher Surgical Trainee; Manchester Royal Infirmary, Oxford Road, Manchester, M13 9WL. ian.farrell@doctors.org.uk

Lyndsay Pearce: BMedSci, BMBS, FRCSEng. Locum Consultant Surgeon, Department of Surgery, Salford Royal NHS Foundation Trust, Stott Lane, Salford, M6 8HD. Lyndsay.pearce@srft.nhs.uk

***Corresponding Author and request for reprints:** Susan J Moug, BSc (Hons), MB ChB, PhD, FRCS. Consultant Surgeon and Honorary Clinical Associate Professor, Department of Surgery, Royal Alexandra Hospital, Paisley, Scotland, PA2 9PN. ++44 141 314 6965. susanmoug@nhs.net

Funding: This study received funding from the Bowel Disease Research Foundation (BDRF) (December 2017). The BDRF were not involved in study design, data collection/ analysis or interpretation of the data. Furthermore, they did not participate in the writing of this work and were not involved of the decision to submit the article for publication. All researchers were independent from the BDRF.

Contributions.

LP and SJM conceived and designed the study in addition to obtaining funding. KLP, JL, JH, JMB, PC, IM, ISF, JH and SJM coordinated the trial. KLP, JL, BC and SJM prepared the first draft of the manuscript and were responsible for the final manuscript. KLP, LP, JL, JMB, PC, IM, ISF, JH and SJM developed the protocol. Local investigators at each site recruited patients. BC authored the statistical analysis plan and carried out the analysis. KLP, JL, BC, JH, LP and SJM interpreted the data. All authors revised manuscript drafts, approved the final manuscript, and contributed intellectually important content. SJM is the guarantor of the paper and takes responsibility for the integrity of the work as a whole, from inception to published article.

ELF Study Group (contributing authors by site in alphabetical order).

| Site | Contributors |
|-------------------------------------|-----------------------|
| Addenbrooke's Hospital | Bryony Ross |
| | Julia Oleksiewicz |
| | Nicola Fearnhead |
| Blackpool Victoria Hospital | Christopher Jump |
| | Jemma Boyle |
| | Alex Shaw |
| | Jonathan Barker |
| Bristol Royal Infirmary | Jane Hughes |
| | Jonathan Randall |
| | Isileli Tonga |
| | James Kynaston |
| | Matthew Boal |
| Countess of Chester Hospital | Nicola Eardley |
| | Elizabeth Kane |
| | Harriet Reader |
| | Sunanda Roy Mahapatra |

| | |
|---------------------------------------|-----------------------|
| | Michael Garner-Jones |
| Croydon University Hospital | Jessica Juliana Tan |
| | Said Mohamed |
| Doncaster Royal Infirmary | Rina George |
| | Ed Whiteman |
| East Cheshire NHS Trust | Kamran Malik |
| | Christopher J Smart |
| | Monica Bogdan |
| East Lancashire Hospitals (Blackburn) | Madhu Parna Chaudhury |
| | Videha Sharma |
| | Daren Subar |
| Furness General Hospital (Barrow) | Panna Patel |
| | Sok-Moi Chok |
| | Evelyn Lim |
| Glan Clywd Hospital | Vedamurthy Adhiyaman |
| | Glesni Davies |
| Glasgow Royal Infirmary | Ellen Ross |
| | Rudra Maitra |
| | Colin W Steele |

| | |
|--|------------------------|
| | Campbell Roxburgh |
| Gloucestershire Royal Hospital | Shelly Griffiths |
| | Natalie S Blencowe |
| | Emily N Kirkham |
| Lorn and Islands Hospital (Oban) | John S Abraham |
| | Kirsty Griffiths |
| Maidstone and Tunbridge Wells NHS Trust | Yasser Abdulaal |
| | Muhammad Rafaih Iqbal |
| | Munir Tarazi |
| Manchester Royal Infirmary | James Hill |
| | Azam Khan |
| | Ian Farrell |
| Mid Essex / Broomfield | Gemma Conn |
| | Jugal Patel |
| | Hyder Reddy |
| Mid Yorkshire NHS Hospitals | Janahan Sarveswaran |
| | Lakshmanan Arunachalam |
| | Afaq Malik |
| | Luca Ponchietti |

| | |
|---|-----------------------|
| Milton Keynes University Hospital NHS Foundation Trust | Krystian Pawelec |
| | Yan Mei Goh |
| | Parveen Vitish-Sharma |
| | Ahmed Saad |
| Musgrove Park (Taunton) | Edward Smyth |
| | Amy Crees |
| | Louise Merker |
| | Nahida Bashir |
| Newport | Gethin Williams |
| | Jennifer Hayes |
| | Kelly Walters |
| | Rhiannon Harries |
| | Rahulpreet Singh |
| Ninewells Hospital (Dundee) | Nikola A Henderson |
| | Francesco M Polignano |
| Queen Alexandra Hospital (Portsmouth) | Ben Knight |
| | Louise Alder |
| | Alexandra Kenchington |
| Queen Elizabeth Hospital (Birmingham) | Yan Li Goh |

| | |
|--|-----------------------|
| | Ilaria Dicurzio |
| | Ewen Griffiths |
| Queen Elizabeth University Hospital (Glasgow) | Ahmed Alani |
| | Katrina Knight |
| | Patrick MacGoey |
| | Guat Shi Ng |
| Royal Albert Edward Infirmary (Wigan) | Naomi Mackenzie |
| | Ishaan Maitra |
| Royal Alexandra Hospital (Paisley) | Susan Moug |
| | Kelly Ong |
| Royal Berkshire Hospital (Reading) | Daniel McGrath |
| | Emanuele Gammeri |
| | Guillame Lafaurie |
| Royal Bolton Hospital | Gemma Faulkner |
| | Gabriele Di Benedetto |
| | Julia McGovern |
| | Bharathi Subramanian |
| Royal Devon and Exeter Hospital NHS | Sunil Kumar Narang |
| | Jennifer Nowers |

| | |
|---|----------------------|
| | Neil J Smart |
| | Ian R Daniels |
| Royal Free Hospital (Hampstead site) | Massimo Varcada |
| | Tanzeela Gala |
| Royal Glamorgan Hospital | Julie Cornish |
| | Zoe Barber |
| Royal Infirmary of Edinburgh | Stephen O'Neill |
| | Richard McGregor |
| | Andrew G Robertson |
| | Simon Paterson-Brown |
| Royal Lancaster Infirmary | Thomas Raymond |
| Royal London Hospital | Mohamed A Thaha |
| | William J English |
| | Cillian T Forde |
| | Heidi Paine |
| | Alpa Morawala |
| Royal Preston Hospital | Ravindra Date |
| | Patrick Casey |
| | Thomas Bolton |

| | |
|--|------------------|
| | Xuan Gleaves |
| | Joshua Fasuyi |
| Royal Surrey County Hospital, Guildford | Sanja Durakovic |
| | Matt Dunstan |
| | Sophie Allen |
| | Angela Riga |
| Salford Royal NHS Foundation Trust | Jonathan Epstein |
| | Lyndsay Pearce |
| | Emily Gaines |
| | Anthony Howe |
| | Halima Choonara |
| Southmead (North Bristol) | Ffion Dewi |
| | Joanne Bennett |
| | Emile King |
| | Kathryn McCarthy |
| Swansea | Greg Taylor |
| | Dean Harris |
| | Hari Nageswaran |
| | Amy Stimpson |

| | |
|---|-------------------|
| Tameside Hospital | Kamran Siddiqui |
| | Lay In Lim |
| University Hospital Crosshouse, Kilmarnock | Christopher Ray |
| | Laura Smith |
| | Gillian McColl |
| University Hospital of South Manchester | Mohammed Rahman |
| | Aaron Kler |
| | Abhi Sharma |
| | Kat Parmar |
| University Hospital of Wales (Cardiff) | Neil Patel |
| | Perry Crofts |
| | Claudio Baldari |
| | Rhys Thomas |
| | Michael Stechman |
| Western General Hospital (Edinburgh) | Roland Aldridge |
| | James O'Kelly |
| | Graeme Wilson |
| Weston general Hospital | Nicholas Gallegos |
| Whiston | Ramya Kalaiselvan |

| | |
|--|--------------------------|
| | Rajasundaram Rajaganesan |
| William Harvey Hospital (East Kent) | Aliya Mackenzie |
| | Prashant Naik |
| | Kaushiki Singh |
| | Harinath Gandraspulli |
| Wirral University Teaching Hospital | Jeremy Wilson |
| | Kate Hancorn |
| | Amir Khawaja |
| | Felix Nicholas |
| | Thomas Marks |
| Wrexham | Cameron Abbott |
| | Susan Chandler |

Mini-abstract

This multicentre study found frailty in 20% of older adults (≥ 65 years) undergoing emergency laparotomy. Independent of age, increasing frailty was associated with a significantly greater risk of 30- and 90-day mortality, in addition to a greater risk of complications and longer length of critical care and overall hospital stay.

Acknowledgements: We gratefully acknowledge support from the North West Surgical Trials Centre (www.nwstc.org.uk). In particular, we would like to thank Laura Marsh and Chris Braithwaite for their help.

Competing interests All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf. All declare funding from the Bowel Disease Research Foundation (BDRF).

Transparency declaration The corresponding author affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Structured Abstract

Objective This study aimed to document the prevalence of frailty in older adults undergoing emergency laparotomy (ELAP) and to explore relationships between frailty and post-operative morbidity and mortality.

Summary Background Data The majority of adults undergoing ELAP are older adults (≥ 65 years) that carry the highest mortality. Improved understanding is urgently needed to allow development of targeted interventions.

Methods An observational multi-centre (n=49) U.K. study was performed (March to June 2017). All older adults undergoing ELAP were included. Pre-operative frailty score was calculated using the progressive Clinical Frailty Score (CFS): 1 (very fit) to 7 (severely frail). Primary outcome measures were the prevalence of frailty (CFS 5-7) and its association to mortality at 90-days post-operative. Secondary outcomes included 30-day mortality and morbidity, length of critical care and overall hospital stay.

Results 937 older adults underwent ELAP: frailty was present in 20%. 90-day mortality was 19.5%. After age and sex adjustment, the risk of 90-day mortality was directly associated with frailty: CFS 5 aOR 3.18 (95%CI 1.24 to 8.14) and CFS 6/7 aOR 6.10 (95%CI 2.26 to 16.45) compared to CFS 1. Similar associations were found for 30-day mortality. Increasing frailty was also associated with: increased risk of complications, length of ICU and overall hospital stay.

Conclusion A fifth of older adults undergoing ELAP are frail. The presence of frailty is associated with greater risks of post-operative mortality and morbidity and is independent of

age. Frailty scoring should be integrated into acute surgical assessment practice to aid decision-making and development of novel post-operative strategies.

INTRODUCTION

Emergency abdominal surgery is performed in every acute hospital in the United Kingdom (U.K.), with the common aim to prevent death and minimise life-altering complications. The majority of these emergency laparotomies are performed on older adults (65 years and older) that when compared to younger adults, have the highest post-operative mortality and morbidity, the highest utilisation of intensive care resources and subsequently, longer lengths of hospital stay.¹⁻⁴ This older adult population has a pre-disposition to more complex clinical needs than younger adults as a consequence of a higher prevalence of multi-morbidity, cognitive impairment, polypharmacy and frailty.^{5,6} However, there is a paucity of good quality research to improve understanding of this expanding older adult population undergoing emergency surgery with many widely used prognostic indicators and tools having been developed from younger, healthier and often elective patient cohorts.⁷⁻¹⁵

Frailty is defined as an objective measure of increased vulnerability and decreased physiological reserve, resulting from the age-associated accumulation of physiological deficits in multiple systems.¹⁶ Frailty results in diminished resilience to physiological insult such as surgery, preventing or impairing recovery and return to pre-existing functional level. In the medical setting where much prior work has focused, routine frailty assessment for older adult admissions has been accompanied by interventions to modify frailty such as the Comprehensive Geriatric Assessment (CGA). The application of the CGA, that can include medical management, physiotherapy, nutrition, pharmacy and occupational health, has led to improved patient survival and an ability to return to pre-admission residence.^{17,18,19} Similar

improved outcomes have been reported in patients with hip fractures as a result of orthogeriatrics input within the first 72 hours of hospital admission.^{20,21} In the emergency general surgery setting, two recent meta-analyses assessing the influence of frailty on older adults documented that the presence of frailty (ranging from 0.5 % to 67.2%) led to significant increases in 30-day mortality, complications and length of hospital stay.^{22,23} However, the inclusion of lower risk elective surgical patients and acute surgical patients managed conservatively, means the prevalence of frailty and its influence on post-operative outcomes in the high-risk emergency laparotomy setting remains unclear.

The primary aims of the Emergency Laparotomy and Frailty (ELF) Study were to determine the prevalence and influence of frailty on 90-day mortality in older adults undergoing emergency laparotomy. The influence of frailty on post-operative outcomes including 30-day mortality, morbidity and length of hospital stay were evaluated as secondary aims.

METHODS

From 20th March to 19th June 2017 all older patients (defined as 65 years and older) undergoing emergency laparotomy at forty-nine registered sites across the U.K. were screened for inclusion into the ELF Study.¹³ This multi-centered prospective cohort study was conceived, designed and led by two established research collaboratives: The North West Research Collaborative (NWRC – www.nwresearch.org) and The Older Persons Surgical Outcomes Collaboration (OPSOC – www.opsoc.eu).

Ethical approval was obtained from National Health Service Research Ethics Committee (Black Country Research Committee: November 2016; 16/WM/0500). The study was registered centrally with the Health Research Authority (England), the NHS Research Scotland Permissions Co-ordinating Centre (Scotland) and the Health and Care Research Permissions Service (Wales). The ELF study was registered online at www.clinicaltrials.gov (NCT02952430).

Site recruitment

Participation from sites across the U.K. was invited using a presentation at the National Research Collaborative meeting in 2016 (www.nationalresearch.org.uk) and the use of social media (@ELFStudy). Fifty-six surgical centers from England, Scotland and Wales registered with data collection received from 49 of those sites.

Patients

The full protocol was published in October 2017 according to STROBE Guidelines.^{24,25} Inclusion criteria were consistent with the established U.K. National Emergency Laparotomy Audit [NELA; www.nela.org.uk/criteria].¹ In general, older patients were included if undergoing an expedited, urgent or emergency surgical abdominal procedure for gastrointestinal pathology (laparoscopic or open procedure) and/ or returning to theatre for any major post-operative complication/dehiscence [Appendix 1]. The classification of Intervention was defined according to the National Confidential Enquire into Patient Outcome and Death (NCEPOD) resulting in elective procedures being excluded.²⁶

Data Collection

Pre-operatively, each older adult had the following recorded: age and gender; number of co-morbidities (allowing calculation of Charlson Co-morbidity index, CCI)^{8,9}; care level/ independence status (selected from home without carers, home with carers, residential home, nursing home, intermediate care, other) and polypharmacy (5 or more current medications). ASA grade (American Society of Anesthesiologists Grade)¹⁰ and P-POSSUM score were calculated as both are routinely collected as part of the NELA.^{11,12} The indication for surgery was documented and in theatre, the procedure performed recorded.

Post-operatively, each patient completed 90-day follow-up. For the first 30 post-operative days, mortality was also recorded in addition to: post-operative complications; length of

critical care stay (level 2 High Dependency Unit, HDU and level 3 Intensive Care Unit, ICU); overall length of hospital stay and readmission to hospital.

Completed anonymous datasets were entered into a specifically designed online secure electronic database (REDCap, www.project-redcap.org) developed and maintained by the North West Surgical Trials Centre (www.nwstc.org.uk).

Frailty Scoring

Each patient's surgical or nursing team assessed frailty pre-operatively as part of standard care. The Clinical Frailty Score (CFS) developed by the Canadian Study of Health and Ageing (CSHA) was used [Appendix 2].²⁷ This 7-point progressive score was developed within an older adult population and is based on clinical judgment with a score of 1 to 4 being classified as non-frail and 5 to 7 as frail. The CFS has been found to be a valid and reproducible score that is simple to understand and apply. OPSOC has used the CFS previously in the emergency surgical population to assess frailty.²⁸

Data Completion and Validation

Regular emails to each registered site and Tweets were used to motivate data collection and update ELF Collaborators. To ensure accurate data collection, data validation was performed on 25% of data fields for 10% of cases at each site after study completion.

Statistical analysis

Sample size justification from previous publication: To detect a 10% difference in 90-day post-operative mortality between frail and non-frail patients (7.5% versus 17.5%), a sample size of 480 was required to maintain 80% power and 5% significance.²⁸

The primary analysis was performed using a multi-level logistic regression of 90-day mortality by frailty, adjusted for age (65–74, and >75 years old) and gender. Each recruiting hospital was fitted as random effects to account for site variation. A secondary analysis of the primary outcome was carried out by presenting the crude odds ratio (OR), and adjusted OR (aOR) with associated 95% CIs and p values. Length of stay was analyzed with a zero inflated negative binomial regression, and presented as the mean increased length of stay (in whole day integers) with associated 95% CI. We carried out an exploratory analysis to investigate the distribution of day 90 mortality, within age and frailty to determine if our findings could be explained in part by patient age. Statistical analyses were carried out using Stata 14 (StataCorp; www.stata.com).

Patient and Public Involvement

There was no patient and public involvement in this study as it was an observational study in a novel research area.

RESULTS

Patients and Procedure

A total of 956 older adults undergoing emergency laparotomy were recruited to the study. Nineteen patients were excluded due to incomplete or incorrectly entered data, leaving 937 patients in the final analysis [Figure 1].

Table 1 displays the pre-operative patient characteristics by frailty score. Overall frailty (CFS 5,6 and 7) was present in 20% of older adults undergoing emergency surgery. The mean age of the patients was 76 years (SD 6.82; range 65 – 99) with over a third (38%) aged 80 or older. Eighty-nine percent of older adults had documented co-morbidity; 54% poly-pharmacy and the majority (83%) were admitted from home without carers. Two thirds of patients (66%) were ASA 3 or more.

The most common indication for surgery was intestinal obstruction (54%). The commonest surgical procedure performed was adhesiolysis (25%) with the majority of surgery performed by an open approach (87%) [Table 2]. These findings are in keeping with the NELA 4th report.¹ Fifty percent of patients had a post-operative complication, with the median length of hospital stay being 13 days (IQR 7-24). 13.7% (123/ 899) of patients required readmission to hospital within 30 days of discharge.

Frailty and 90-day Post-operative Mortality

Overall 90-day mortality for all patients was 19.5% (183/930). As the frailty score increased the risk of 90-day mortality increased: patients scoring CFS 6 and 7 had the highest risk of 90-day mortality (OR 5.89, 95% CI 2.19 to 15.86; $p=0.001$) compared to patients scoring CFS1 [Table 3 and Figure 2]. After accounting for patient age and sex, increasing frailty was still associated with increased risk of 90-day mortality: aOR for patients who were mildly frail (CFS 5) and moderately/ severely frail (CFS 6/7) was 3.18 (95%CI 1.24 to 8.14; $p=0.016$) and 6.10 (95%CI 2.26 to 16.45; $p<0.001$) respectively compared to CFS 1 (very fit).

Frailty and Secondary Outcomes

Overall 30-day mortality for all patients was 14.6% (137/ 937) and a similar association with frailty was seen for 30-day mortality as for 90-day mortality: CFS 5 aOR 9.79 (95%CI 2.23 to 42.91; $p=0.002$) and for CFS 6/7 aOR 10.40 (95%CI 2.24 to 48.18; $p=0.003$) [Table 3]. Increasing frailty score was associated with increased risk of post-operative complications: aOR for CFS 5 was 4.56 (95% CI 2.17 to 9.60; $p=0.001$) and 3.92 (95%CI 0.35 to 4.19; $p=0.001$) for CFS 6/7 compared to those older adults scoring CFS 1.

There was increased length of post-operative hospital stay and stay in ICU in frail patients [Table 3]. For hospital stay: aOR for CFS 5 was 1.44 (95% CI 1.10 to 1.89; $p=0.008$) and 1.62 (1.19 to 2.20; $p=0.002$) for CFS 6/7 compared to adults scoring CFS 1. For ICU length of stay: CFS 5 aOR of 2.15 95%CI 1.15 to 3.96; $p=0.02$) and CFS 6/7 aOR 4.18 (95% CI 2.11 to 8.03; $p<0.001$)

compared to CFS 1. Frail patients stayed in ICU for 2 days (IQR 1-4) versus 1 day (IQR 0-3) for non-frail. There was no association with frailty to the length of time in HDU or to 30-day re-admission.

Frailty, by Age mortality distribution

Tables 4A and 4B show that frailty is distributed throughout the range of ages and not restricted to the very old age patient groups (85+). Observation of this distribution shows that patients with a higher frailty score had a higher mortality rate. The marginal 90-day mortality rates were increased from 11.1% to 50% for CFS of 1 to 7 and the marginal mortality rate for increased patient age was 15.9% to 25.3% for patients aged 65-70, compared to those aged 85-90 [Table 4A]. Similar results were found for the 30-day mortality showing that frailty results in a wider distribution of predicted mortality than age [Table 4B]. In addition when compared to P-POSSUM, it can be seen that frailty stratifies the older adult into a greater number of prognostic groups and also allows risk-prediction of 30-day morbidity through the range of scoring, whilst P-POSSUM does not [Table 5].

DISCUSSION

This study is the first to prospectively document the prevalence of frailty in older adults undergoing emergency laparotomy. Frailty was present in a fifth of patients pre-operatively placing them at significantly greater risk of 30- and 90-day mortality. In addition, as the frailty score increased, so did the risk of post-operative complications, and the length of stay in intensive care and in hospital overall. With frailty found to be independent of age, this work supports integration of frailty scoring to all older adults admitted as a surgical emergency to guide peri-operative strategies for this high-risk complex population.

There are several strengths to this work. First, it focused solely on a large population of older patients undergoing emergency laparotomy improving understanding of this overlooked group that carry the highest risk for post-operative death. Second, the data was collected prospectively and was multi-centred minimising geographical bias that could occur with local or regional studies. Third, validation of a proportion of each site's data optimised data completion. Finally, the Clinical Frailty Score (CFS) appears to be a simple tool to use with only a small percentage of frailty scores not entered despite the substantial over-recruitment.

The authors cannot exclude selection bias where a local recruiter may have not contributed a consecutive series of older patients undergoing emergency laparotomy. This is likely to reflect the pragmatic nature of this work. In addition, only a small proportion of patients undergoing

surgery had the highest frailty score (CFS 7), perhaps as a consequence of being unfit for surgery. To overcome this, CFS 6 was combined with CFS 7 for statistical analysis where appropriate.

Previous work in the emergency surgical setting has found frailty to be associated with a greater risk of mortality and longer hospital stays.^{28,29,30,31} However, direct comparisons to this current work are difficult due to their limitations of: retrospective analysis; single center only; inclusion of non-operatively managed older adult patients and/ or small patient numbers. In a recently published prospective multi-centered U.K. study (n=2,279) assessing all patients admitted to an acute surgical unit, frailty was found to exist in all age groups and was not exclusive to the older adult population.³² This study included patients regardless of whether an operation was performed (62% had non-operative management), however the finding of frailty to be independent of age with an almost linear relationship to 90-day mortality is consistent with our study. Furthermore, with our results also showing incremental increases in frailty scores predict greater risks of post-operative complications and longer hospital stays, we believe that the CFS should be interpreted as an individual score rather than the traditional binary classification of frail or not frail.

Clinicians have long been interested in prognostic scores resulting in many being available. All have their limitations in the older adult emergency laparotomy setting including: predicting mortality only (APACHE II, ASA); multiple variables making calculation labor intensive (APACHE

II); not validated in older adult populations (Charlson, ASA); being based on diagnosed co-morbidity with no attempt to differentiate well-controlled morbidity from severe functionally limiting morbidity (Charlson) or requiring intra-operative details to complete score (P-POSSUM).⁷⁻¹² Indeed, NELA and The Royal College of Surgeons of England have acknowledged and circumvented such limitations by using P-POSSUM or an equivalent to define a high-risk patient to trigger specific process pathways rather than to determine prognosis.^{1,15} The Clinical Frailty Score in comparison predicts both morbidity and mortality, is straightforward to calculate, is validated in older adult surgical populations and can be determined pre-operatively.

Implications for clinicians and policymakers

Frailty scoring should be performed pre-operatively on all older adults undergoing an emergency laparotomy to aid with the complex decision-making and peri-operative care. Few of the already established prognostic scores are easy for the patient and their family to understand, which is potentially detrimental to shared decision-making.^{33,34} In contrast, frailty is a concept of which many members of the public may be aware. This underlying knowledge may provide a platform for improved understanding with patients and their family members when discussing not only their operative risks of dying, but of having significant life-altering complications and a prolonged and difficult recovery.

Once the decision for surgery is made, the frailty score could lead targeted peri-operative pathways. For example, patients with CFS 4 to 7 have the highest risk of mortality where early

involvement and assessment by critical care would optimise post-operative planning. Treatment over the first few days post-operatively is focused on continuing treatment for the initial pathology (e.g. sepsis, renal support, ventilator support, wound management) led by critical care and surgeons. This could be improved by working alongside geriatricians and the CGA with the key aim of minimising any complications from frailty.^{1,6,15,35} For example: protecting muscle mass (early mobility or movement whether on or off a ventilator); maintaining respiratory capacity (timed regular physiotherapy); nutrition and energy balance (parental or enteral). In contrast, a patient CFS 3 might only spend a short time in critical care before continuing their targeted rehabilitation prescribed by a hospitalist or geriatrician, but led by the surgical ward nurses. This modified emergency laparotomy CGA creates an opportunity for targeted training within both surgical and geriatric curriculums to allow multidisciplinary peri-operative care.^{6,36}

CONCLUSION

Frailty is present in 20% of older adults undergoing emergency laparotomy and is independent of age. As frailty increases, the older adult is at greater risk of post-operative mortality and morbidity. These findings support the integration of pre-operative frailty assessment and identify the urgent need to develop novel post-operative strategies to improve outcomes for this complex high-risk group of health service users.

References

1. The National Emergency Laparotomy Audit (NELA). www.nela.org.uk

2. Desserud KF, Veen T, Søreide K. Emergency general surgery in the geriatric patient. *Br J Surg*. 2016;103:e52-61.
3. Svenningsen P, Manoharan T, Foss NB, Lauritsen ML, Bay-Nielsen M. Increased mortality in the elderly after emergency abdominal surgery. *Dan Med J*. 2014;61:A4876.
4. Chow WB, Merkow RP, Cohen ME, Bilimoria KY, Ko CY. Association between postoperative complications and reoperation for patients undergoing geriatric surgery and the effect of reoperation on mortality. *Am Surg*. 2012;78:1137-1142.
5. Moug SJ, Stechman M, McCarthy K, Pearce L, Myint PK, Hewitt J; Older Persons Surgical Outcomes Collaboration. Frailty and cognitive impairment: Unique challenges in the older emergency surgical patient. *Ann R Coll Surg Engl*. 2016 Mar;98:165-169.
6. Pearce L, Bunni J, McCarthy K, Hewitt J. Surgery in the Older Person: Training needs for the provision of multidisciplinary care. *Ann R Coll Surg Eng*. 2016; 98:367- 370.
7. Ozban M, Birsen O, Senel M, Ozden A, Kabay B. The analysis of scoring systems predicting mortality in geriatric emergency abdominal surgery. *Ulus Travma Acil Cerrahi Derg*. 2015;21:182-1826.
8. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol*. 1994;47:1245–1251.
9. St-Louis E, Iqbal S, Feldman LS, Sudarshan M, Deckelbaum DL, Razek TS, et al. Using the age-adjusted Charlson comorbidity index to predict outcomes in emergency general surgery. *J Trauma Acute Care Surg*. 2015;78:318-323.
10. Saklad M. Grading of patients for surgical procedures. *Anesthesiology*. 1941;2:281–284.

11. Copeland GP, Jones D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg.* 1991;78:355–360.
12. Prytherch DR, Whiteley MS, Higgins B, Weaver PC, Prout WG, Powell SJ. POSSUM and Portsmouth POSSUM for predicting mortality. *Br J Surg.* 1998;85:1217–1220.
13. Office for National Statistics. What does the 2011 Census tell us about older people? 6th September 2013.
<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/ageing/articles/whatdoesthe2011censustellusaboutolderpeople/2013-09-06>
14. World Population Prospects. The 2017 Revision. Key Findings and Advance Tables. United Nations Department of Economic and Social Affairs Population Division. New York 2017.
https://esa.un.org/unpd/wpp/Publications/Files/WPP2017_KeyFindings.pdf
15. The High-Risk general Surgical Patient: Raising the Standard. The Royal College of Surgeons of England Updated Recommendations on the Perioperative Care of the High-Risk General Surgical Patient, 2018. <https://www.rcseng.ac.uk/news-and-events/media-centre/press-releases/high-risk-general-surgical/>
16. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet.* 2013;381:752-762.
17. Wells JL, Seabrook JA, Stolee P, Borrie MJ, Knoefel F. State of the art in geriatric rehabilitation. Part I: review of frailty and comprehensive geriatric assessment. *Arch Phys Med Rehabil.* 2003;84:890-897.

18. Gill TM, Baker DI, Gottschalk M, Gahbauer EA, Charpentier PA, de Regt PT, et al. A prehabilitation program for physically frail community-living older persons. *Arch Phys Med Rehabil.* 2003;84:394-404.
19. Deschodt M, Flamaing J, Haentjens P, Boonen S, Milisen K. Impact of geriatric consultation teams on clinical outcome in acute hospitals: a systematic review and meta-analysis. *BMC Med.* 2013;11:48.
20. Mangram AJ, Mitchell CD, Shifflette VK, Lorenzo M, Truitt MS, Goel A, et al. Geriatric trauma service: a one-year experience. *J Trauma Acute Care Surg.* 2012;72:119-22.
21. Mangram AJ, Shifflette VK, Mitchell CD, Johnson VA, Lorenzo M, Truitt MS, et al. The creation of a geriatric trauma unit "G-60". *Am Surg.* 2011;77:1144-1146.
22. Sandini M, Pinotti E, Persico I, Picone D, Bellelli G, Gianotti L. Systematic review and meta-analysis of frailty as a predictor of morbidity and mortality after major abdominal surgery. *BJS Open.* 2017 1: 128-137.
23. Hewitt J, Long S, Carter B, Bach S, McCarthy K, Clegg A. The prevalence of frailty and its association with clinical outcomes in general surgery: a systematic review and meta-analysis. *Age Ageing.* 2018 1;47(6):793-800.
24. Parmar KL, Pearce L, Farrell I, Hewitt J, Moug S. Influence of frailty in older patients undergoing emergency laparotomy: a UK-based observational study. *BMJ Open.* 2017;7:e017928.
25. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet.* 2007;370:1453-7.

26. Elective and Emergency Surgery in the Elderly: An Age old Problem 2010. National Confidential Enquiry into Patient Outcome and Death. www.ncepod.org.uk
27. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173:489-495.
28. Hewitt J, Moug SJ, Middleton M, Chakrabarti M, Stechman MJ, McCarthy K, et al. Prevalence of frailty and its association with mortality in general surgery. *Am J Surg*. 2015;209:254-259.
29. Farhat JA, Velanovich V, Falvo AJ et al. Are the frail destined to fail. Frailty index as predictor of surgical morbidity and mortality in the elderly. *J Trauma Acute Care Surg*. 2012, 71:1526-31. Discussion 30-31.
30. Bellal J, Bardiya Z, Viraj P et al. Emergency General Surgery in the Elderly: Too Old or Too frail? *J Am Coll Surg*. 2016; 222:805-813.
31. Orouji Jokar T, Ibraheem K, Rhee P et al. Emergency general surgery specific frailty index: a validation study. *J Trauma Acute Care Surg*. 2016; 81:254-260.
32. Hewitt J, Carter B, McCarthy K et al. frailty predicts mortality in all emergency surgical admissions regardless of age. An observational study. *Age and Ageing* 2019 0:1-7.
33. Smith AK, Williams BA, Lo B. Discussing overall prognosis with the very elderly. *N Engl J Med*. 2011;365:2149-2151.
34. Realising Realistic Medicine. Catherine Calderwood, Chief Medical Officer, Scotland. <https://beta.gov.scot/news/realising-realistic-medicine/>

35. Oliver DM, Bassett MG, Poulton TE, Anderson ID, Grocott MP, Moonesinghe SR. Organisational factors and mortality after an emergency laparotomy: multi-level analysis of 39 903 National Emergency Laparotomy Audit patients. *BJA*. 2018;1346-1356.
36. Barbas AS, Haney JC, Henry BV, Heflin MT, Lagoo SA. Development and implementation of a formalized geriatric surgery curriculum for general surgery residents. *Gerontol Geriatr Educ*. 2014;35:380-394.

Figure 1: ELF Study flow diagram.

Table 1: Characteristics of Older Adults undergoing Emergency Laparotomy, by Frailty score.

Table 2: Indication and type of surgical procedure performed on Older Adults undergoing emergency laparotomy.

Table 3: Frailty and Primary and Secondary outcomes in Older Adults undergoing Emergency Laparotomy.

Figure 2: Scatterplot displaying the relationship between Frailty and 90-day mortality in Older Adults undergoing Emergency laparotomy.

Table 4A: The distribution of 90-Day mortality within Frailty by age in Older Adults undergoing Emergency Laparotomy.

Table 4B: The distribution of 30-Day mortality within Frailty by age in Older Adults undergoing Emergency Laparotomy.

Table 5: Comparison of Frailty status to P-Possum by Actual 30-day Mortality and Morbidity.

Appendix 1: National Emergency Laparotomy Audit (NELA) Inclusion and Exclusion Criteria [1].

Appendix 2: The Clinical Frailty Score (CFS) developed by the Canadian Study of Health and Ageing (CSHA).